

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/53, 9/02, C12P 7/22, 7/02	A1	(11) International Publication Number: WO 96/14419
		(43) International Publication Date: 17 May 1996 (17.05.96)

(21) International Application Number: PCT/GB95/02588

(22) International Filing Date: 2 November 1995 (02.11.95)

(30) Priority Data:
9422205.6 3 November 1994 (03.11.94) GB(71) Applicant (for all designated States except US): BRITISH
GAS PLC [GB/GB]; Rivermill House, 152 Grosvenor Road,
London SW1V 3JL (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): FLITSCH, Sabine, Lahja
[GB/GB]; 14 West Saville Terrace, Edinburgh EH9 3DZ
(GB). NICKERSON, Darren, Paul [CA/CA]; Holywell
Manor, Manor Road, Oxford OX1 3UH (GB). WONG,
Luet-Lok [GB/GB]; 29 Crotch Crescent, Oxford OX3 0JL
(GB).(74) Agent: MORGAN, David, J.; British Gas plc, Intellectual
Property Dept., 59 Bryanston Street, London W1A 2AZ
(GB).(81) Designated States: AU, CN, CZ, JP, KR, NZ, PL, RU, SG,
SK, UA, US, European patent (AT, BE, CH, DE, DK, ES,
FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).**Published***With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*(54) Title: MUTANT MONO-OXYGENASE CYTOCHROME P-450_{cam}

(57) Abstract

A mutant of the mono-oxygenase cytochrome P-450_{cam} in which the tyrosine residue at position 96 and/or the cysteine residue at position 334 is replaced by the residue of any amino acid except phenylalanine.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Mutant mono-oxygenase cytochrome P-450_{cam}

The present invention relates to a mutant of the mono-oxygenase cytochrome P-450_{cam} and method of oxidising certain organic compounds with the mutant.

Mono-oxygenases catalyse the selective oxidation of non-functionalised hydrocarbons using oxygen¹, and are therefore of great interest for potential use in organic synthesis. However, progress in this area has been hampered by the difficulty in isolating sufficient quantities of enzyme and the associated electron-transfer proteins. Despite the availability of amino acid sequences of more than 150 different cytochrome P-450 mono-oxygenases, to date structural data of only three are available^{2,3,4}, and few have been successfully over-expressed in bacterial systems⁵.

One cytochrome P-450 mono-oxygenase, which is soluble and can be expressed in sufficient quantities, is the highly specific P-450_{cam} from *P.putida* which catalyses the regio- and stereo-selective hydroxylation of camphor (1) to 5-exo-hydroxycamphor⁶. The high resolution crystal structure of P-450_{cam} has been determined², and since the mechanism of action of this bacterial enzyme is believed to be very similar to that of its mammalian counterparts, it has been used as a framework on which models of mammalian enzymes are based.

The nucleotide sequence and corresponding amino acid sequence of P-450_{cam} have been described⁵. The location of an active site of the enzyme is known and structure-function relationships have been investigated^{13,14}. Mutants of P-450_{cam} have been described, at the 101 and 185 and 247 positions¹⁵, and at the 87 position¹⁶. A mutant in which tyrosine 96 has been changed to phenyl alanine-96 has been described^{12,17,18}. But in all these cases the papers report effects of the mutations on the mechanisms of known oxidation reactions. There is no teaching or suggestion that mutation might be used to provide biocatalysts for oxidation of different substrates.

In an attempt to find new biocatalysts, we have initiated a project which aims to redesign the active site of P-450_{cam}, such that it is able to carry out specific oxidations of organic molecules which are not substrates for the wild-type protein. Our initial aim was to incorporate an "aromatic pocket" into the P-450_{cam} active site, which would encourage the binding of substrates containing aromatic side-chains.

In addition, a surface residue remote from the active site was identified (cysteine-334) with effects on protein handling and stability. The cysteine is responsible for unwanted dimerisation of the protein during purification and an alanine residue was therefore substituted for the cysteine in order to improve both of these properties.

The three dimensional structure of P-450_{cam} shows the active site to provide close van der Waals contact with the hydrophobic groups of camphor as shown in Figure 1. The aromatic residues (Y96, F87 and F98) are grouped together and line the substrate binding pocket, with a hydrogen bond between tyrosine 96 and the camphor carbonyl oxygen maintaining the substrate in the correct orientation to ensure the regio- and stereo-specificity of the reaction. Replacement of any of these aromatic residues with a smaller, hydrophobic non-aromatic side-chain could provide the desired "aromatic pocket".

Molecular modelling was used to investigate the likely effects of point mutations to the three aromatic residues. The program GRID⁷ was used to calculate an energy of interaction between an aromatic probe and possible mutants of cytochrome P-450_{cam} where these residues were changed to alanine (F87A, Y96A and F98A). The results were then examined graphically using the molecular modelling package Quanta⁸.

The mutant F98A appeared to have the strongest binding interaction within the active site cavity accessible to the aromatic probe, with that of Y96A being slightly smaller, and that of F87A being substantially less. It was decided in the first instance to mutate tyrosine 96 to alanine as it is more central to the binding pocket, whereas phenylalanine 98 is in a groove to one side. Also, removal of tyrosine 96 should

decrease the specificity of the enzyme towards camphor due to the loss of hydrogen bonding to the substrate.

According to one aspect of the present invention a mutant of the mono-oxygenase cytochrome P-450_{cam} is provided in which the tyrosine residue at position 96 and/or the cysteine residue at position 334 is replaced by the residue of any amino acid except phenylalanine.

According to another aspect of the present invention a mutant of the mono-oxygenase cytochrome P-450_{cam} is provided in which the tyrosine residue at position 96 and/or the cysteine residue at position 334 is replaced by another amino acid residue, which mutant has the property of catalysing the oxidation of any one of the following:- polycyclic aromatic hydrocarbons, linear or branched alkanes, diphenyl and biphenyl compounds including halogenated variants of such compounds and halogenated hydrocarbons.

According to yet another aspect of the present invention a method is provided of oxidising a compound selected from a polycyclic aromatic hydrocarbon, a linear or branched alkane, a diphenyl or biphenyl compound including a halogenated variant of such a compound or a halogenated hydrocarbon, the method comprising contacting the selected one of the compounds under oxidising conditions with mono-oxygenase cytochrome P-450_{cam} in which the tyrosine residue at position 96 and/or the

cysteine residue at position 334 is replaced by another amino acid residue.

Preferably the amino acid is selected from any one of the following:- alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, proline, serine, threonine, tryptophan, tyrosine and valine except that in the case of the tyrosine residue at position 96, the amino acid is not tyrosine and in the case of the cysteine residue at position 334, the amino acid is not cysteine.

The amino acid which replaces tyrosine at position 96 is conveniently one of the small hydrophobic amino acids, e.g. alanine, glycine, valine, leucine or isoleucine, with alanine being preferred as exemplified below.

Alternatively the amino acid replacing tyrosine at position 96 may be one of the charged amino acids, e.g. a negatively charged acid such as aspartic acid or glutamic acid for hydrogen bonding to a positively charged substrate; or a positively charged compound such as lysine, arginine or histidine for hydrogen bonding to a negatively charged substrate which are not members of the camphor family.

The mutation at position 96 is believed to be the key which enables the mutant enzymes to catalyse the oxidation of a relatively wide range of organic substrates. Other amino

acids adjacent to the active site of the enzyme may also be mutated in order to change the shape and specificity of the active site. These other amino acids include those at positions 87, 98, 185, 244, 247, 295 and 297. It is envisaged that the amino acid at one or more of these positions may be replaced by: a small hydrophobic amino acid so as to enlarge the active site; or a large hydrophobic amino acid so as to reduce the size of the active site; or by an amino acid having an aromatic ring to bond to a corresponding aromatic ring of a substrate.

Regarding the oxidising reactions, the conditions are described in the literature references attached. The enzyme system typically includes putidaredoxin and putidaredoxin reductase together with NADH as co-factors in addition to the mutant enzyme. Various classes of organic compounds are envisaged:-

- i) The organic compound is an aromatic compound, either a hydrocarbon or a compound used under conditions in which it does not inactivate or denature the enzyme. Since the mutation has been effected with a view to creating an aromatic-binding pocket in the surface of the enzyme, the mutant enzyme is capable of catalysing oxidation of a wide variety of aromatic compounds. Oxidation of example aromatic and polyaromatic compounds is demonstrated in the experimental section below and is believed very

surprising given that the wild-type enzyme catalyses the oxidation of only members of the camphor family.

- ii) The organic compound may be a hydrocarbon, e.g. aliphatic or alicyclic, carrying a functional group. An aromatic protective group is added to the functional group prior to the oxidation reaction and removed from the functional group after the oxidation reaction. A suitable aromatic group is a phenyl group. The aromatic protection group is expected to hold the substrate in place. Thus the protecting group serves two purposes: firstly it makes the substrate more hydrophobic and hence increases binding to the hydrophobic enzyme pocket; secondly it holds the substrate in place at the active site. Thus, with the correct aromatic protection group, both regio- and stereo-selective hydroxylation of the substrate may be achieved. The example of cyclohexylbenzene is described in the experimental section below.

Examples of monofunctionalised hydrocarbons are cyclohexyl, cyclopentyl and alkyl derivatives (Scheme 1). The oxidation products of these compounds are valuable starting materials for organic synthesis, particularly when produced in a homochiral form. A range of aromatic protecting groups are envisaged, e.g. benzyl or naphthyl ethers and benzoyl or naphthoyl esters and amides (Scheme 1). Of interest are also benzoxazole groups as carboxyl protecting groups and N-benzyl oxazolidine groups as

ald hyde protecting groups. Both can be easily cleaved after the enzymatic oxidation and hav previously b en described in the literature for the microbial oxidations of aldehydes and acids.

- iii) The organic compound is a C5 to C12 aliphatic or alicyclic hydrocarbon. Oxidation of cyclohexane and linear hydrocarbons is demonstrated in the experimental section below and once again it is believed quite surprising given that the wild-type enzyme catalyses the oxidation of only members of the camphor family.
- iv) The organic compound is a halogenated aliphatic or alicyclic hydrocarbon. Oxidation of lindane (hexachlorocyclohexane) is also described below.

Reference is directed to the accompanying drawings in which Figure 2 is a gas chromatograph of diphenylmethane (A) and hydroxylated product formed following incubation with P-450_{cam} Y96A mutant.

Based on the above considerations, mutant proteins were constructed which contained alanine, lysine, valine, or phenylalanine instead of tyrosine at position 96 (Y96). Additional mutants were constructed in which these active site replacements were combined with the surface mutation of cysteine at position 334 to alanine. Lastly several active site mutations and the surface mutation were combined in one

protein to constitute a multiple mutant enzyme. The genes encoding cytochrome P-450_{cam}, and its natural electron-transfer partners puridaredoxin and putidaredoxin reductase, were amplified from the total cellular DNA of *P. putida* using the polymerase chain reaction (PCR). The expression vector/*E. coli* host combinations employed were pRH1091⁹ in strain JM109 for P-450_{cam}, pUC 118 in strain JM109 for putidaredoxin, and pGLW11 in strain DH5^α for putidaredoxin reductase. Oligonucleotide-directed site-specific mutagenesis was carried out using an M13mp19 subclone by the method of Zoller and Smith¹⁰, and mutant selection was by the method of Kunkel¹¹.

The mutant Y96A was shown to catalyse the hydroxylation of camphor (1), although compared to the wild-type enzyme the reaction was less selective, similar to that reported for the mutant Y96F¹². This decrease in selectivity can be attributed to the loss of the hydrogen bond between Y96 and camphor. The properties of wild-type and Y96A proteins were further investigated with a variety of binding and activity assays.

Binding of potential substrates was investigated by spectroscopic methods. The wild-type enzyme in the absence of substrate is in the 6-co-ordinated, low-spin form with a weakly bound water occupying the sixth co-ordination site, and shows a characteristic Soret maximum at 391 nm. Binding of the substrate analogues adamantanone (2), adamantane (3) and norbornane (4) also fully converted the haem to the high-spin

form. However, diphenylmethane (5) did not give a shift in the absorption spectrum.

The Y96A mutant, while giving the same results for compounds (3) and (4), was not fully converted to the high-spin form even when (1) and (2) were added in excess. Most interestingly however, and in contrast to the wild-type, Y96A showed partial conversion to the haem to the high-speed form with diphenylmethane, indicating binding of this compound to the mutant protein.

As expected, the dissociation constants (K_{app}) for camphor and adamantanone are increased in Y96A. On the other hand, the K_{app} values for the hydrophobic substrates adamantane and norbornane are reduced, indicating that the enzyme pocket has become more selective for hydrophobic substrates. The greatest change in binding was obtained with diphenylmethane, which bound poorly to wild-type protein, but showed greatly enhanced affinity for the Y96A mutant (Table 1).

Once binding of diphenylmethane by the Y96A protein had been established, catalytic substrate turnover was investigated. The mutant protein was reconstituted with putidaredoxin and putidaredoxin reductase. Diphenylmethane (5) was added and the mixture was incubated with NADH and oxygen.

A solution containing 10 μ M putidaredoxin, 2 μ M putidaredoxin reductase, 1 μ M cytochrome P-450_{cam} mono-oxygenase (wild-type

or mutant) and 1 mM diphenylmethane in 100 mM KCl, 20 mM KH_2PO_4 , pH 7.4 was preincubated at 25°C in a shaker for 5 min. The enzymatic reaction was initiated by firstly adding NADH to a total concentration of 2 mM. Further four aliquots of NADH (to increase the NADH concentration by 1 mM each time) were added in intervals of 5 min and the reaction quenched after 30 min by adding 0.5 ml chloroform. The chloroform layer was analysed by gas chromatography.

Organic extracts of the crude incubation mixture were analysed by gas chromatography. Only one major new peak was detected by GC (see Figure 2), which had the same retention time as an authentic sample of para-hydroxydiphenylmethane (6). The other aromatic hydroxylation products, the ortho and meta isomers, had different retention times. Further confirmation of the identity of the product as structure (6) was provided by mass spectrometry, which gave the correct mass peak at 184.

Using the above experimental techniques, the inventors have investigated a considerable number of organic compounds as substrates for both the wild-type P-450_{cam} enzyme and also the mutant version Y96A. Further work has included mutants designated Y96V; Y96L; Y96F; C334A; the combined mutant F87A, Y96G, F193A and the combined active site and surface mutants of Y96A, C334A; Y96V, C334A; Y96L, C334A; Y96F, C334A; F87A, Y96G, F193A, C334A.

The results for Y96A are set out in Table 2, in which structurally related molecules are grouped together. Those substrates where oxidation has been demonstrated by means of NADH turnover are marked with a + sign.

Spin high/low: numbers show the percentage of P-450 (OD_{417} 0.2-0.4) converted from the low- to high-spin equilibrium state in the presence of 200 μ M test compound, in phosphate buffer (40 mM phosphate, 68 mM potassium, pH 7.4). Spin state equilibrium is assessed with a UV/vis spectrophotometer: low spin at OD_{417} and high spin at OD_{392} nd; not done.


Vs DTT: numbers show the percentage displacement of DTT (200 μ M) bound to P-450 by competition with test compounds (200 μ M) in phosphate buffer. DTT binding to P-450 results in absorbance peaks at OD_{374} and OD_{461} , so displacement is measured with a UV/vis spectrophotometer.

Examples are included in Table 2(a) to 2(h) for each class of compounds identified in points i) to iv) above.

Reaction products for some substrate compounds have been purified by high performance liquid chromatography and identified by mass spectroscopy, nuclear magnetic resonance, and/or co-elution. Table 3 details the NADH consumption for oxidation of small linear, branched and cyclic hydrocarbons by the mutant Y96A, C334A. Table 4(a) to 4(h) details the






product distributions for mutant and substrat combinations
where this has been elucidated to date.

Scheme 1:



15

Table 1:

		K_{app} (μ M) ^a	
		WT	Y96A
	<u>1</u>	6.3	12
	<u>2</u>	12	28
	<u>3</u>	8.4	1.4
	<u>4</u>	330	92
	<u>5</u>	>1500 ^b	73

^a Values are the average of two independent measurements using the method of Sligar (S.G. Sligar, *Biochemistry*, **1976**, *15*, 5399 - 5406). The value of K_{app} is strongly dependent on the concentration of K^+ in the buffer. At $[K^+] > 150$ mM, K_{app} for camphor is 0.6 μ M for both wildtype and Y96A. Data in this table were determined at $[K^+] = 70$ mM in phosphate buffer, pH 7.4, in order to avoid salting out of substrates at higher ion concentrations.

^b Saturation not reached.

SUBSTITUTE SHEET (RULE 26)

Table 2(a)

P450cam-substrate interactions		Wild type		Mutant Y96A		Wild type		Mutant Y96A	
Subgroup: 1-ring		Δ Spin high/low	Vs DTT	Δ Spin high/low	Vs DTT	NADH turnover?	GC?	NADH turnover?	GC?
Benzene		-	-	-	-				
Toluene		-	-	30	30				
Ethylbenzene		-	-	40	40				
Styrene		-	-	30	30				
Cyclohexene		-	5	40	40				
1,3-Cyclohexadiene		nd	nd	nd	nd				
1,4-Cyclohexadiene		-	5	15	20				
Cyclohexane		-	-	60	60			+	
Hexane		-	-	70	60			+	
Methylcyclohexane		50	50	100	70				
(S)-(+)-Carvone		10	60	10	80				

SUBSTITUTE SHEET (RULE 26)

Table 2(b)











P450cam-substrate interactions		Wild type		Mutant Y96A		Wild type		Mutant Y96A	
Subgroup: 2-ring, Naphthalene		Δ Spin high/low	Vs DTT	Δ Spin high/low	Vs DTT	NADH turnover?	GC?	NADH turnover?	GC?
		-	-	15	20				
1-Ethyl-naphthalene		-	-	5	20				
									
2-Ethyl-naphthalene		-	-	10	20				
									
2-Naphthylacetate		-	5	-	5				
									
1-Naphthylacetate		-	5	-	5				
									
1-Naphthylpropionate		-	20	0	20				
									
1-Naphthylbutyrate		-	5	-	5				
									
Naphthylphenylketone		-	5	-	5				
									
1,2-Dihydronaphthalene		5	20	30	90				
									
1,2,3,4-Tetrahydro naphthalene		5	10	40	40				
									

Table 2(c)













p450cam-substrate interactions		Wild type		Mutant Y96A		Wild type		Mutant Y96A	
Subgroup: 2-ring, DPM		Δ Spin high/low	Vs DTT	Δ Spin high/low	Vs DTT	NADH turnover?	GC?	NADH turnover?	GC?
	Diphenylmethane	-	5	45	nd			+	+
	Diphenylether	10	5	20	50				
	Benzophenone	-	20	-	20				
	Cyclohexylphenylketone	-	30	60	nd				
	Phenylbenzoate	-	5	-	-				
	N-Phenylbenzylamine	-	5	45	nd				
	Bibenzyl	-	-	55	55				
	cis-Stilbene	-	20	40	50				
	Biphenyl	-	20	-	90				
	Cyclohexylbenzene	20	20	80	nd				
	trans-Stilbene	-	-	-	-				
	Benzylether	-	5	55	nd				

Table 2(d)









P450cam-substrate Interactions		Wild type		Mutant Y96A		Wild type		Mutant Y96A	
		Δ Spin high/low	Vs DTT	Δ Spin high/low	Vs DTT	NADH turnover?	GC?	NADH turnover?	GC?
Subgroup: 3-ring									
	Anthracene	.	.	20	20
	Phenanthrene	.	.	20	20	.	.	+	.
	Fluorene	.	.	.	50
	2-Fluorencarboxaldehyde	.	.	.	50
	9-Fluorenone	.	20	.	5
	Anthrone	.	5	.	5
	Anthraquinone
	2-Ethylanthraquinone

Table 2 (e)






P450cam-substrate interactions	Wild type		Mutant Y96A		Wild type		Mutant Y96A	
	Δ Spin high/low	Vs DTT	Δ Spin high/low	Vs DTT	NADH turnover?	GC?	NADH turnover?	GC?
Subgroup: 4,5-ring								
 Chrysene	-	-	-	-				
 1,2-Benzanthracene	-	-	-	-				
 Fluoranthene	-	5	20	10				
 Pyrene*	-	-	-	-				
 Perylene*	-	-	-	-				

Table 2(f)





P450cam-substrate interactions		Wild type		Mutant Y96A		Wild type		Mutant Y96A	
Subgroup: Cyclic Alkanes		Δ Spin high/low	Vs DTT	Δ Spin high/low	Vs DTT	NADH turnover?	GC?	NADH turnover?	GC?
 α -Decahydronaphthalene		nd	nd	nd	nd				
	 <i>trans</i> -Decahydro naphthalene	20	10	90	70				
 Cyclohexane		.	.	60	60			+	
 Methylcyclohexane		50	50	100	70				

Table 2 (g)



P450cam-substrate interactions		Wild type		Mutant Y96A		Wild type		Mutant Y96A	
Subgroup: n-Alkanes		Δ Spin high/low	Vs DTT	Δ Spin high/low	Vs DTT	NADH turnover?	GC?	NADH turnover?	GC?
n-Pentane		-	5	55	40			+	
n-Hexane		-	-	60	40			+	
n-Heptane		5	5	60	40			+	
n-Octane		-	5	80	45			+	
n-Nonane		-	-	70	45			+	
n-Decane		nd	nd	nd	nd				
n-Undecane		nd	nd	20	20				
n-Dodecane		nd	nd	5	5				
$\text{CH}_3(\text{CH}_2)_{14}\text{CH}_3$	n-Hexadecane	-	-	-	-				
$\text{CH}_3(\text{CH}_2)_{15}\text{CH}_3$	n-Heptadecane	-	-	-	-				
$\text{CH}_3(\text{CH}_2)_{11}\text{OSO}_3\text{Na}$	SDS	-	20	-	60				
$\text{CH}_3(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_7\text{CO}_2\text{H}$	Oleic acid*	-	10?	-	20?				
$[(\text{CH}_3)_2\text{CH}(\text{CH}_2)_3\text{CH}(\text{CH}_3)(\text{CH}_2)_3\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2]_2$									
	Squalene	-	-	-	20				
	Isoprene	-	-	10	10				

Table 2(h)




P450cam-substrate Interactions		Wild type		Mutant Y96A		Wild type		Mutant Y96A	
		Δ Spin high/low	V_s DTT	Δ Spin high/low	V_s DTT	NADH turnover?	GC?	NADH turnover?	GC?
Subgroup: Camphor-like		80	80	80	80				
	(1R)-(-)-Camphorquinone								
		40	70	50	80				
	(1R)-(-)-Fenchone								
		50	80	90	90				
	Dicyclopentadiene								

Table 3

Turnover of Small Alkanes by P450cam Mutants

All mutants listed below also contain the C334A mutation.

Turnover rate measured as NADH consumption rate (nmole NADH/nmole P450cam/s).





Alkane substrate: Main chain length	Name	Wild type	Y96A
C4	<i>n</i> -butane	-	-
C4	2-methyl butane	background	4.6
C4	2,3-dimethyl butane	background	16.8
C4	2,2-dimethyl butane	background	14.0
C5	<i>n</i> -pentane	background	5.8
C5	2-methyl pentane	3.8	11.7
C5	3-methyl pentane	1.3	14.2
C5	2,4-dimethyl pentane	0.2	12.6
C5	2,2-dimethyl pentane	5.2	12.8
C5	2,2,4-trimethyl pentane	0.9	5.3
C5	3-ethyl pentane	background	16.2
C6	<i>n</i> -hexane	background	6.0
C6	2-methyl hexane	background	10.6
C7	<i>n</i> -heptane	2.7	4.4
C7	2-methyl heptane	background	2.1
C7	4-methyl heptane	1.4	10.2
C8	<i>n</i> -octane	background	5.8
C7	cycloheptane	4.4	42.5

Product structures and distributions following oxidation of substrates with P450cam active site mutants.

"background" - typical background NADH oxidation rate is 0.07 nmole NADH (nmole P450cam)⁻¹ sec⁻¹

Table 4(a)

Product structure and distributions following oxidation of substrates with P450cam active site mutants. All mutants shown below also contain the C334A mutation.

Cyclohexylbenzene		Products (%) for mutants:				
Products	WT	Y96A	Y96F	Y96L	Y96V	
 3-ol D or L	43	20	54	38	28	
 3-ol " " L or D	20	20	27	23	39	
 Trans -4-ol	25	15	6	23	10	
 Cis- 4-ol	12	45	13	16	23	
Total products(area/10 ³)	0.8	7.4	1.1	10.4	12.5	






Table 4(b)

Phenylcyclohexene		Products (%) for mutants:	
Products		WT	Y96A
	3-one (A)	24	25
	3-ol (B)	76	75
Total products(area/10 ⁶)		42	36








Table 4(c)

Naphthalene Products	Products (%) for mutants:									
	WT	Y96A	Y96F	Y96L	Y96V	F87A-F96G-	F193A			
1-ol	100	100	100	100	100	100	100			
2-ol	0	0	0	0	0	0	0			
Total products (area/10 ⁵)	(0.016)	1.1	2.4	0.7	1.4	0.1	0.1			





 P450cam
 Mutants




Table 4(d)										
Phenanthrene Products		WT	Y96A	Y96F	Y96L	Y96V	F87A-F96G-F193A	Products (%) for mutants:		
A		38	49	41	35.5	41	27			
B		15	23	31	41	38	41			
C		12	13	5	9	11	3			
D		35	15	23	14.5	10	29			
Total products (area/10 ⁶)		0.075	7.0	4.5	2.8	1.6	0.065			




Table 4(e)

Fluoranthene Products	Products (%) for mutants:					
	WT	Y96A	Y96F	Y96L	Y96V	F87A-F96G- F193A
A	0	84	-	-	-	0
B	0	16	-	-	-	100
Total products	0	2.7	-	-	-	0.2
(area/10 ⁶)						




Table 4(f)

Pyrene Products		Products (%) for mutants:							
	WT	Y96A	Y96F	Y96L	Y96V	F87A-F96G-F193A			
A	0	40	43	23	30	33			
B	0	43.6	29	64.5	55	40			
C	0	5	12.5	7.9	12	20			
D	0	11.4	15.5	4.6	3	7			
Total products		0	1.2	1.5	1.5	1.6	0.02		
		(area/10 ⁶)							




Table 4(g)

**Lindane Products
(hexachlorocyclohexane)**

	Products (%) for mutants	
	WT	Y96A


100

100

A

43.5

7.5

Turnover rate
nmole NADH (nmoleP450)⁻¹s⁻¹

Hexachlorocyclohexane

Table 4(h)

Hexane Products	Products (%) for mutants:	
	Y96F	Y96A
2-hexanone	10	15
3-hexanone	16	28
2-hexanol	24	26
3-hexanol	50	32
Relative activity (WT = 1)	18.2	25.5

2-Methyl hexane Products	Products (%) for mutants:	
	Y96F	Y96A
2-methyl-2-hexanol	72	74
5-methyl-2-hexanone	16	14
2-methyl-3-hexanol	7	4
5-methyl-2-hexanol	5	8
Relative activity (WT = 1)	2.3	2.6

REFERENCES

1. "Cytochrome P-450: Structure, Mechanism, and Biochemistry", ed. P R Ortiz de Montellano, Plenum Press, New York, 1986.
2. T L Poulos, B C Finzel and A J Howard, J. Mol. Biol., 1987, 195, 687-700.
3. J A Peterason, U.-Y. Lu, J Geisselsoder, S Graham-Lorence, C Carmona, F Witney, and M C Lorence, J. Biol. Chem., 1992, 267, 14193-14203.
4. K G Ravichandran, S S Boddupali, C A Hasemann, J A Peterson, and J Deisenhofer, Science, 1993, 261, 731-736.
5. B P Unger, I C Gunsalus, and S G Sligar, J. Biol. Chem., 1986, 261, 1158-1163; J S Miles, A W Munro, B N Rospendowski, W E Smith, J McKnight, and A J Thomson, Biochem. J., 1992, 288, 503-509; T H Richardson, M J Hsu, T Kronbach, H J Barnes, G Chan, M R Waterman, B Kemper, and E F Johnson, Arch. Biochem. Biophys., 1993, 300, 510-516; S S Boddupalli, T Oster, R W Estabrook, and J A Peterson, J Biol. Chem., 1992, 267, 10375-10380; H Li K Darish and T L Poulos, J Biol. Chem., 1991, 266-11909-11914.

6. I C Gunsalus and G C Wagner, *Methods Enzym* 1., 1978, 52, 166-188.
7. P J Goodford, *J Med. Chem.*, 1985, 28, 849-857.
8. Quanta 4.0, Molecular Simulations Inc., 16 New England Executive Park, Burlington, MA 01803-5297.
9. J E Baldwin J M Blackburn, R J Heath, and J D Sutherland, *Bioorg, Med. Chem. Letts.* 1992, 2, 663-668.
10. M J Zoller and M Smith, *Nucleic Acids Res.*, 1982, 10, 6487.
11. T A Kunkel, *Proc. Natl. Acad. Sci.*, 1985, 82, 488-492.
12. C Di Primo, G Hui Bin Hoa, P. Douzou, and S Sligar, *J. Biol. Chem.*, 1990, 265, 5361-5363.
13. D Filipovic, *Biochemical and Biophysical Research Communications*, Vol. 189, No. 1, 1992, November 30, 1992, pages 488-495.
14. S G Sligar, D Filipovic, and P S Stayton, *Methods in Enzymology*, Vol. 206, pages 31-49.
15. P J Loida and S G Sligar, *Protein Engineering*, Vol. 6, No. 2, pages 207-212, 1993.

16. S F Tuck t al., Th Journal of Biological Chemistry, Vol. 268, No. 1, January 5, 1993, pages 269-275.
17. W M Atkins and S G Sligar, The Journal of Biological Chemistry, Vol. 263, No. 35, 15 December 1988, pages 18842-18849.
18. W M Atkins and S G Sligar, Biochemistry 1990, 29, 1271-1275.

CLAIMS

1. A mutant of the mono-oxygenase cytochrome P-450_{cam} in which the tyrosine residue at position 96 and/or the cysteine residue at position 334 is replaced by the residue of any amino acid except phenylalanine.
2. A mutant of the mono-oxygenase cytochrome P-450_{cam} in which the tyrosine residue at position 96 and/or the cysteine residue at position 334 is replaced by another amino acid residue, which mutant has the property of catalysing the oxidation of any one of the following:- polycyclic aromatic hydrocarbons, linear or branched alkanes, diphenyl and biphenyl compounds including halogenated variants of such compounds and halogenated hydrocarbons.
3. A mutant as claimed in claim 1 or claim 2 in which the amino acid is selected from any one of the following:- alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, proline, serine, threonine, tryptophan, tyrosine and valine except that in the case of the tyrosine residue at position 96, the amino acid is not tyrosine and in the case of the cysteine residue at position 334, the amino acid is not cysteine.


4. A mutant as claimed in any of claims 1 to 3 in which the amino acid residue at one or more of the positions 87, 98, 185, 244, 247, 295 and 297 is replaced by another amino acid residue.
5. A method of oxidising a compound selected from a polycyclic aromatic hydrocarbon, a linear or branched alkane, a diphenyl or biphenyl compound including a halogenated variant of such a compound or a halogenated hydrocarbon, the method comprising contacting the selected one of the compounds under oxidising conditions with mono-oxygenase cytochrome P-450_{cam} in which the tyrosine residue at position 96 and/or the cysteine residue at position 334 is replaced by another amino acid residue.
6. A method as claimed in claim 5 in which the amino acid is selected from any one of the following:- alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, proline, serine, threonine, tryptophan, tyrosine and valine except that in the case of the tyrosine residue at position 96, the amino acid is not tyrosine and in the case of the cysteine residue at position 334, the amino acid is not cysteine.
7. A method as claimed in claim 5 or claim 6 in which the amino acid residue at one or more of the positions 87,

98, 185, 244, 247, 295 and 297 is replaced by another amino acid residue.


8. A mutant of the mono-oxygenase cytochrome P-450_{cam} substantially as hereinbefore described with reference to the accompanying drawings and/or examples.
9. A method of oxidising a compound selected from a polycyclic aromatic hydrocarbon, a linear or branched alkane, a diphenyl or biphenyl compound including a halogenated variant of such a compound or a halogenated hydrocarbon substantially as hereinbefore described with reference to the accompanying drawings and/or examples.

1/2

FIG.1.



2/2



INTERNATIONAL SEARCH REPORT

Inter-Office Application No
PCT/95/02588

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/53 C12N9/02 C12P7/22 C12P7/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 263, no. 35, 15 December 1988 pages 18842-18849, W.M. ATKINS ET AL. 'The role of active site hydrogen bonding in cytochrome P-450 cam as revealed by site-directed mutagenesis.' cited in the application	2,8
Y	see the whole document ---	1,3-7
X	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 265, no. 10, 5 April 1990 pages 5361-5363, C. DI PRIMO ET AL. 'Mutagenesis of a single hydrogen bond in cytochrome p450 alters cation binding and heme solvation.' cited in the application	2,8
Y	see the whole document ---	1,3,4
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

27 March 1996

Date of mailing of the international search report

22.04.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hix, R

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BIOCHEMISTRY, vol. 29, no. 5, 6 February 1990 pages 1271-1275, W. M. ATKINS ET AL. 'Tyrosine-96 as a natural spectroscopic probe of the cytochrome P-450cam active site' cited in the application	2,8
Y	see the whole document ---	1,3,4
Y	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 1, 5 January 1993 pages 269-275, S. F. TUCK ET AL. 'Active sites of the cytochrome p450cam {CYP101} F87W and F87A mutants.' cited in the application see the whole document ---	1,3-7
E	WO,A,95 34679 (US HEALTH ;GONZALEZ FRANK J (US); IDLE JEFFREY R (GB)) 21 December 1995 see the whole document ---	1-9
P,A	WO,A,95 16041 (CIBA GEIGY AG ;ROYAL VETERINARY & AGRICULTURA (DK); KOCH BIRGIT MA) 15 June 1995 see the whole document ---	1-9
A	PSYCHIATRIC GENETICS 4 (4). 1994. 215-218. ISSN: 0955-8829, DAWSON E ET AL 'An association study of debrisoquine hydroxylase (CYP2D6) polymorphisms in schizophrenia.' see the whole document ---	1-9
A	THE LANCET, vol. 339, no. 8806, 6 June 1992 pages 1375-1377, C.A.D. SMITH ET AL 'Debrisoquine hydroxylase gene polymorphism and susceptibility to Parkinson's disease.' see the whole document -----	1-9

INTERNATIONAL SEARCH REPORT

Information on family members

International Application No

PCT/GB 02588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9534679	21-12-95	AU-B- 2860295	05-01-96
WO-A-9516041	15-06-95	AU-B- 1108295	27-06-95

Form PCT/ISA/210 (patent family annex) (July 1992)

THIS PAGE BLANK (USPTO)